



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A novel role for the mineralocorticoid receptor in glucocorticoid driven vascular calcification

Citation for published version:

Zhu, D, Rashdan, NA, Chapman, KE, Hadoke, PWF & MacRae, VE 2016, 'A novel role for the mineralocorticoid receptor in glucocorticoid driven vascular calcification', *Vascular pharmacology*, vol. 86, pp. 87-93. <https://doi.org/10.1016/j.vph.2016.04.005>

Digital Object Identifier (DOI):

[10.1016/j.vph.2016.04.005](https://doi.org/10.1016/j.vph.2016.04.005)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Vascular pharmacology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Accepted Manuscript


A novel role for the mineralocorticoid receptor in glucocorticoid driven vascular calcification

Dongxing Zhu, Nabil A. Rashdan, Karen E. Chapman, P.W.F. Hadoke, Vicky E. MacRae

PII: S1537-1891(15)30124-5
DOI: doi: [10.1016/j.vph.2016.04.005](https://doi.org/10.1016/j.vph.2016.04.005)
Reference: VPH 6311

To appear in: *Vascular Pharmacology*

Received date: 29 December 2015
Revised date: 22 March 2016
Accepted date: 18 April 2016



Please cite this article as: Zhu, Dongxing, Rashdan, Nabil A., Chapman, Karen E., Hadoke, P.W.F., MacRae, Vicky E., A novel role for the mineralocorticoid receptor in glucocorticoid driven vascular calcification, *Vascular Pharmacology* (2016), doi: [10.1016/j.vph.2016.04.005](https://doi.org/10.1016/j.vph.2016.04.005)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**A Novel Role for the Mineralocorticoid Receptor in
Glucocorticoid Driven Vascular Calcification.**

Dongxing Zhu¹, Nabil A Rashdan¹, Karen E. Chapman², P. W. F. Hadoke², Vicky E.
MacRae¹

1: The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of
Edinburgh, Midlothian, EH25 9RG, UK

2: University/BHF Centre for Cardiovascular Sciences, University of Edinburgh, The
Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh, EH16 4TJ, UK

**Address for Correspondence*

Dr Vicky MacRae,
The Roslin Institute and Royal (Dick) School of Veterinary Studies,
University of Edinburgh,
Easter Bush,
Roslin,
Midlothian. EH25 9RG.
UK

Tel 00 44 (0) 131 651 9156

Email vicky.macrae@roslin.ed.ac.uk

Abstract

Vascular calcification, which is common in the elderly and in patients with atherosclerosis, diabetes and chronic renal disease, increases the risk of cardiovascular morbidity and mortality. It is a complex, active and highly regulated cellular process that resembles physiological bone formation. It has previously been established that pharmacological doses of glucocorticoids facilitate arterial calcification. However, the consequences for vascular calcification of endogenous glucocorticoid elevation have yet to be established. Glucocorticoids (cortisol, corticosterone) are released from the adrenal gland, but can also be generated within cells from 11-keto metabolites of glucocorticoids (cortisone, 11-dehydrocorticosterone [11-DHC]) by the enzyme, 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1). In the current study we hypothesised that endogenous glucocorticoids facilitate vascular smooth muscle cell (VSMC) calcification and investigated the receptor-mediated mechanism underpinning this process.

In vitro studies revealed increased phosphate-induced calcification in mouse VSMCs following treatment for 7 days with corticosterone (100 nM; 7.98 fold; $P<0.01$), 11-DHC (100 nM; 7.14 fold; $P<0.05$) and dexamethasone (10 nM; 7.16 fold; $P<0.05$), a synthetic glucocorticoid used as a positive control. Inhibition of 11 β -HSD isoenzymes by 10 μ M carbenoxolone reduced the calcification induced by 11-DHC (0.37 fold compared to treatment with 11-DHC alone; $P<0.05$). The glucocorticoid receptor (GR) antagonist mifepristone (10 μ M) had no effect on VSMC calcification in response to corticosterone or 11-DHC. In contrast, the mineralocorticoid receptor (MR) antagonist eplerenone (10 μ M) significantly decreased corticosterone- (0.81 fold compared to treatment with corticosterone alone; $P<0.01$) and 11-DHC- (0.64 fold compared to treatment with 11-DHC alone; $P<0.01$) driven VSMC calcification, suggesting this glucocorticoid effect is MR-driven and not GR-driven. Neither corticosterone nor 11-DHC altered the mRNA levels of the osteogenic markers *Pit-1*, *Osx* and *Bmp2*. However, DAPI staining of pyknotic nuclei and flow cytometry analysis of surface Annexin V expression showed that corticosterone induced apoptosis in VSMCs.

This study suggests that in mouse VSMCs, corticosterone acts through the MR to induce pro-calcification effects, and identifies 11 β -HSD-inhibition as a novel potential treatment for vascular calcification.

Introduction

Vascular calcification is a marker of increased cardiovascular disease risk in ageing, including in diabetes, atherosclerosis and chronic kidney disease (CKD) (Mackenzie and MacRae, 2011; Zhu et al., 2012). The etiology of mineral accumulation within the vasculature shares many similarities with that of bone formation. Indeed, several studies have reported that vascular smooth muscle cells (VSMCs), the predominant cell type involved in vascular calcification, can undergo phenotypic transition to osteoblastic, chondrocytic and osteocytic cells in a calcified environment (Johnson et al., 2005; Zhu et al., 2011). Furthermore, phosphate accelerates this trans-differentiation process, with the loss of characteristic smooth muscle markers and the increased expression of osteoblastic markers (eg. Osterix, Pit-1 and BMP2) (Montes de Oca et al., 2014; Nakagawa et al., 2010; Zhu et al., 2013). Vascular calcification can also proceed through mechanisms involving the reciprocal loss of recognised calcification inhibitors including inorganic pyrophosphate (PPi), fetuin A and Matrix Gla Protein (Johnson et al., 2005; Mackenzie et al., 2012; Murshed et al., 2005; Reynolds et al., 2005).

Physiological glucocorticoids - primarily cortisol in humans and corticosterone in rats and mice - are steroid hormones produced by the adrenal cortex. Local glucocorticoid action on target tissues is determined by intracellular metabolism by the two isozymes of 11β -hydroxysteroid dehydrogenase (11β -HSD) which catalyze interconversion of active cortisol and corticosterone with inert cortisone and 11-dehydrocorticosterone (Hadoke et al., 2013). 11β -HSD type 1, a predominant reductase in most intact cells, catalyzes the regeneration of active glucocorticoids, thus amplifying cellular action. 11β -HSD2 is a high-affinity dehydrogenase that inactivates glucocorticoids (Chapman et al., 2013). Both isozymes of 11HSD are modestly expressed in the blood vessel wall, suggesting that they can influence vascular function by regulating local availability of active glucocorticoids (Hadoke et al., 2013).

In the absence of 11β -HSD2, endogenous glucocorticoids can bind to the mineralocorticoid receptor (MR) as well as the glucocorticoid receptor (GR) (Richardson et al., 2015). Both MR and GR belong to the same nuclear hormone receptor superfamily, and share high sequence identity. MR has higher affinity for glucocorticoids than GR, and both receptors are expressed in the cells of the vasculature (Hadoke et al., 2006). Glucocorticoids can activate

the MR in VSMCs (11 β -HSD2 is not expressed here), inducing pathways that are central to cell proliferation and differentiation (Molnar et al., 2008).

Glucocorticoids are frequently permissive, co-operative or synergistic (Wada, 2008). Indeed a permissive role of glucocorticoids in triggering cell transdifferentiation has previously been established in the conversion of pancreatic cells into hepatocytes (Shen et al., 2000). Furthermore, glucocorticoids exert complex actions on calcium mobilisation and bone metabolism, regulating bone resorption and formation (Hartmann et al., 2016), intestinal calcium absorption and renal calcium excretion (Diaz de Barboza et al., 2015). Therefore it is essential to establish the consequences for vascular calcification of endogenous glucocorticoid elevation given the high circulating calcium levels commonly observed in patients with this pathology (Block, 2000).

Dexamethasone, a potent synthetic glucocorticoid which is primarily active at the GR, induces an osteoblastic differentiation pathway in many different mesenchymal-derived cell types *in vitro* (Bellows et al., 1990; Cheng et al., 1994; Walsh et al., 2001), including VSMCs (Mori et al., 1999; Shui & Scutt, 2002). Whilst the established pro-calcification actions of dexamethasone on VSMCs (Kirton et al., 2006; Mori et al., 1999) are presumed to be mediated via GR, plausibly endogenous corticosteroids may modulate VSMC calcification via MR. This is therapeutically important to ascertain, as vascular calcification is independently correlated with adverse cardiac events (Zhu et al., 2012), and MR antagonism is highly successful in reducing mortality in heart failure (Vizzardi et al., 2014); aldosterone antagonists such as spironolactone and eplerenone have been shown to improve cardiovascular outcomes and prevent ischaemic events in cardiovascular patients (Pitt et al., 1999; 2003). Corticosterone has been shown to induce rapid MR signalling in VSMCs that involves mitogen-activated protein kinase kinase (MEK)/extracellular signal-regulated kinase (ERK)-dependent pathways, suggesting that glucocorticoids may contribute to vascular disease via MR receptor signalling (Molnar et al., 2008). Recent studies have shown that aldosterone-induced activation of MR promotes osteoblastic differentiation and calcification of VSMCs (Jaffe et al., 2007) through a mechanism involving the stimulation of spironolactone-sensitive, PiT-1 dependent signalling (Voelkl et al., 2013).

An additional level of control over endogenous corticosteroid action is provided by the HSD isoenzymes, whose role in vascular calcification has yet to be elucidated. The induction of

local glucocorticoid generation through increased 11β -HSD1 expression (>10 fold) and activity (>4 fold) by inflammatory cytokines and glucocorticoids is well documented in fibroblasts and osteoblasts (Kaur et al., 2010; Sun & Myatt, 2003), which both have the capacity to calcify (Staines et al., 2014; Boraldi et al., 2014).

It is therefore essential to establish the consequences for vascular calcification of endogenous glucocorticoid elevation and potential strategies for inhibition of calcification. The aims of this study were to undertake *in vitro* murine VSMC calcification studies to investigate both the identity of the receptor and the role of the 11β -HSD isoenzymes in corticosterone-induced calcification.

Materials and Methods

Mice

All animal experiments were performed under UK Home Office licensed approval in accordance with Directive 2010/63/EU of the European Parliament and were maintained in accordance with Home Office guidelines for the care and use of laboratory animals. C57BL/6 mice were supplied by Charles River Laboratories (Harlow, Essex, UK).

Preparation of VSMCs

Mice were euthanized by cervical dislocation. Primary murine VSMCs were isolated as described (Mackenzie et al., 2011). Briefly, after removal of the adventitia, the aorta was opened to expose the endothelial layer under a dissection microscope. Tissues from eight animals were pooled and incubated with 1mg ml^{-1} trypsin (Invitrogen, Paisley, UK) for 10 min in order to enable the removal of any remaining adventitia and endothelium through further dissection. Following overnight incubation at 37°C in a humidified atmosphere of 95% air/5% CO_2 in “growth medium” (α -MEM supplemented with 10% Fetal Bovine Serum and 1% gentamicin, all from Invitrogen), tissues were digested with 425U/ml collagenase type II (Worthington Biochemical Corporation, Lakewood, USA) for 5 h. Cell suspensions were centrifuged at 2000 g for 5 min. The cell pellet was washed and resuspended in growth medium. Isolated VSMCs were passaged in growth medium twice in T25 tissue culture flasks (Greiner Bio-one, GmbH, Frickenhausen, Baden-Wurttemberg, Germany) coated with $0.25\mu\text{g/cm}^2$ laminin (Sigma, Poole, UK) to promote maintenance of the contractile

differentiation state (Johnson et al., 2008). VSMCs were subsequently seeded at a density of $1.5 \times 10^4/\text{cm}^2$ in 12-well plates.

Induction of VSMC calcification

In vitro calcification of VSMCs was induced by culturing cells in growth medium containing 3 mM inorganic phosphate (a mixture of NaH_2PO_4 and Na_2HPO_4 , pH 7.4, Sigma) for up to 14 days, with a medium change every 3 days, as previously described (Zhu et al., 2015; 2016). The effects of glucocorticoids in FBS were assessed through comparison of charcoal-stripped and standard FBS (Life Technologies Ltd). Cells were treated with corticosterone (1-100 nM) (Sigma), 11-DHC (1-100 nM) (Steraloids, Newport, USA), carbenoxolone (10 μM) (Sigma), dexamethasone (1-100 nM) (Sigma), mifepristone (10 μM) (Sigma) or eplerenone (10 μM) (Sigma). The *in vitro* levels of corticosterone and 11-DHC used in the present study reflect those found *in vivo*. Plasma corticosterone levels in mice range from 20 nM (nadir, morning) to ~150 nM (peak, evening) and stress levels are typically 200-250 nM. Basal levels of plasma 11-DHC in mice have been reported at 2-5 nM and stress levels >30 nM (Harris et al., 2001).

Determination of VSMC calcification

Calcium deposition was quantified by HCl leaching, as described previously (Zhu et al., 2014). Briefly, cells were washed twice with phosphate buffered saline (PBS) and incubated with 0.6M HCl at room temperature for 24 h. Calcium content was determined colorimetrically by a stable interaction with phenolsulphonethalein using a commercially available kit (Randox Laboratories Ltd., County Antrim, UK), corrected for total protein concentration (Bio-Rad Laboratories Ltd, Hemel Hempstead, UK), and presented as a fold change compared with control. Calcium deposition was also evaluated by alizarin red staining. Cells were washed twice with PBS, fixed in 4% paraformaldehyde for 5 min at 4°C , stained with 2% alizarin red (pH 4.2) for 5 min at room temperature and rinsed with distilled water.

Analysis of gene expression

VSMCs were treated with corticosterone (Sigma) or 11-DHC (Steraloids, Newport, USA), for 48 h in serum free α -MEM (Invitrogen). RNA was extracted using RNeasy total RNA (Qiagen Ltd, Crawley, West Sussex, UK), according to the manufacturer's instructions. RNA was quantified and reverse transcribed as previously described (Mackenzie et al., 2014).

Levels of specific mRNAs were measured using the SYBR green detection method (Roche, East Sussex, UK) as previously reported (Staines et al., 2014). Primers were obtained from Qiagen (sequences not disclosed) for *PiT-1* (NM_001159593), *Bmp2* (NM_007553) and *Msx2* (NM_013601).

Quantification of apoptosis

On reaching confluence, cells were serum starved for 24 h, then treated with 100 nM corticosterone for 48 h. Cells were harvested by trypsinization and re-suspended in 25 µl 1% trypan blue (diluted 50% in PBS). Live cells, which exclude trypan blue and dead cells (stained blue) were counted using a haemocytometer, and the results expressed as the percentage of cells that were dead. Apoptotic VSMCs were determined by manually counting pyknotic nuclei after staining with DAPI (Invitrogen) as previously described (Guicciardi et al., 2011). Additionally, cells in different stages of apoptosis were analysed by flow cytometry using the TACS Annexin-V-FITC apoptosis detection kit (R&D systems, Abingdon, UK), according to the manufacturer's instructions. Non-apoptotic cells do not stain with either Annexin-V FITC or propidium iodide. Early apoptotic cells are stained with Annexin-V FITC but not propidium iodide (green fluorescence). Late apoptotic cells are stained with both Annexin-V FITC and propidium iodide (dual green and red fluorescence). Necrotic cells are only stained with propidium iodide (red fluorescence). 10,000 cell events were recorded on a BD FACS Calibur and data were analyzed with FlowJo 8.8.4 flow cytometry analysis software (Tree Star Inc., Ashland, Oregon, USA).

Statistical analysis

General Linear Model analysis and the Students t-test were used to assess the data. All data are expressed as the mean +/- S.E.M. Statistical analysis was performed using Minitab 16. $P < 0.05$ was considered to be significant.

Results

Glucocorticoids facilitates VSMC calcification

We initially examined the effects of physiological glucocorticoids on the calcification of VSMCs, with the synthetic glucocorticoid dexamethasone used as a positive control. Since arterial calcification is highly correlated with elevated serum Pi levels, VSMCs were cultured in growth medium containing high (3 mM) Pi as previously described (Zhu et al.,

2015;2016). Cells were treated with dexamethasone, corticosterone or 11-DHC (1-100 nM) for up to 7 days. Dexamethasone treatment significantly increased calcium deposition of VSMCs (17.16 fold, $P<0.05$, Fig. 1A). Both corticosterone and 11-DHC significantly increased calcium deposition in VSMCs cultured with un-stripped FBS (1.51 fold, $P<0.001$ and 1.72 fold respectively, $P<0.001$; Fig. 1B). Surprisingly, 11-DHC was more potent than corticosterone, with a significant effect at 10 nM, compared to 100 nM for corticosterone. Using FBS stripped of steroids markedly increased the magnitude of the effects of both corticosterone and 11-DHC on VSMC calcification (7.98 fold, $P<0.01$ and 7.14 fold, $P<0.05$ respectively; Fig. 1C). The 11 β -HSD inhibitor carbenoxolone notably reduced 11-DHC-induced calcification of VSMCs (0.37 fold compared to treatment with 11-DHC alone, $P<0.001$, Fig. 1D), confirming the 11 β -HSD isoenzymes as key regulators of intracellular glucocorticoid levels (Chapman et al., 2013).

Corticosterone and 11-DHC facilitate VSMC calcification through the MR.

In order to establish whether corticosterone acts via GR or MR to induce VSMC calcification, antagonists of both MR (eplerenone) and GR (mifepristone) were employed. Mifepristone (10 μ M) itself potentiated VSMC calcification, but did not alter the response to corticosterone (Fig. 2A) or 11-DHC (Fig. 2B). However, eplerenone (10 μ M) significantly attenuated the effect of corticosterone (0.81 fold compared to treatment with corticosterone alone; $P<0.01$) and 11-DHC- (0.64 fold compared to treatment with 11-DHC alone; $P<0.01$) on VSMC calcification.

Corticosterone and 11-DHC induce VSMC apoptosis.

Published data implicate the transdifferentiation of VSMCs to an osteoblast-like phenotype as a mechanism underlying the effects of dexamethasone upon vascular calcification (Mori et al., 1999). We therefore next tested whether the same is true for physiological glucocorticoids. Interestingly, neither corticosterone nor 11-DHC altered mRNA levels of key osteogenic markers: *Pit-1* (Fig. 3A), *Osx* (Fig. 3B) and *Bmp2* (Fig. 3C). Recent reports have highlighted apoptosis as essential for the initiation and progression of phosphate-induced vascular calcification (Son et al., 2008). Therefore we undertook a detailed assessment of apoptosis following glucocorticoid treatment. Corticosterone treatment significantly reduced cell viability (2.07 fold; $P<0.01$; Fig. 4A), and conversely increased cell death (2.53 fold; $P<0.05$; Fig. 4B) and apoptosis (Fig. 4C-E) as determined by DAPI staining of pykrotic nuclei and FACS analysis of cells positively stained for Annexin V.

Discussion

It is well established that glucocorticoids mediate changes in vascular growth, function and structure (Hadoke et al., 2006). Previous studies in VSMCs have reported the pro-calcification effects of dexamethasone, a potent synthetic glucocorticoid. Here we demonstrate for the first time the facilitation of vascular calcification by both the active physiological glucocorticoid corticosterone and the inactive metabolite 11-DHC.

Currently 11 β -HSD inhibition is therapeutically employed in topical preparations for the management of mouth ulcers (Hirata et al., 2013). Furthermore, 11 β -HSD-inhibition has been shown to correct insulin resistance in rodent models of CKD, a hallmark of this disease in patients being vascular calcification (Block et al., 2000). The *in vitro* experiments undertaken in this study therefore identify 11 β -HSD inhibition as a plausible treatment for vascular calcification, and requires further interrogation *in vivo*.

Whilst VSMC calcification is facilitated by GR selective dexamethasone (Fig. 1; Mori et al., 1999), and macrophage-specific GR inactivation reduces vascular calcification in a mouse model of atherosclerosis (Preusch et al., 2008), the pro-calcification effects of corticosterone and 11-DHC do not appear to be mediated through GR signalling. Furthermore, whilst dexamethasone has been previously reported to accelerate the osteogenic differentiation of vascular pericytes (Kirton et al., 2006) and bovine VSMCs (Mori et al., 1999), in the present study *Osterix*, *Bmp2* and *PiT-1* expression were unaltered in VSMCs treated with corticosterone or 11DHC. Interestingly, these data contrast with a previous demonstration of up-regulated *PiT-1* in response to MR signalling (Voelkl et al., 2013). This paradigm likely reflects different mechanisms underpinning the calcification permitted by synthetic and physiological glucocorticoids. Indeed, our studies suggest that the pro-calcification effects of corticosterone and 11-DHC are mediated directly through MR, corroborating previous work in VSMCs showing the activation of MR signalling by glucocorticoids (Molnar et al., 2008). Furthermore, our data complement previous work revealing that CVCs (calcifying vascular cells; subpopulations of VSMCs which have been found to spontaneously calcify *in vitro*) contain MRs which function as transcriptional regulators that can be activated by both aldosterone and cortisol (Jaffe et al., 2007). Indeed aldosterone administration has been shown to directly stimulate CVC calcification, an effect abolished by aldosterone antagonism with spironolactone (Jaffe et al., 2007). Moreover, vascular calcification facilitated by

hyperaldosteronism due to klotho deficiency has also been shown to be mitigated by spironolactone treatment in mice (Voelkl et al., 2013). Further studies are therefore required to directly compare the effects of aldosterone treatment on VSMC calcification with that of corticosterone and 11-DHC.

The present study also highlights that the effect of corticosterone and 11-DHC on driving VSMC calcification is more pronounced when charcoal stripped media is used to remove endogenous steroid ligands, suggesting an activating effect of endogenous MR ligands in normal, unstripped serum. These data support comparable findings demonstrating that aldosterone-induced CVC calcification via MR activation is also enhanced using charcoal stripped serum (Jaffe et al., 2007).

Mechanistically, our data reveal that corticosterone reduces cell viability and stimulates VSMC apoptosis. This process is essential for the initiation and progression of phosphate-induced vascular calcification (Son et al., 2008), with apoptotic bodies exposing phosphatidylserine on the outer membranes, generating a potential calcium-binding site suitable for hydroxyapatite deposition (Proudfoot et al., 2000; Skrtic & Eanes, 1992). These results support previous reports demonstrating that glucocorticoids inhibit VSMC proliferation (Longenecker et al., 1984; Versaci et al., 2002) and induce apoptosis in a range of cell types including neuronal cells, growth plate chondrocytes and thymocytes (Chmielewski et al., 2000; Chrysis et al., 2005; Liu et al., 2011 and Li et al., 2014).

In conclusion, we have undertaken *in vitro* murine VSMC studies to provide new insights into the role of physiological glucocorticoids in vascular calcification. Our study suggests that corticosterone acts through the MR to induce pro-calcification effects. This data may have important health ramifications for patients receiving MR blockers. The previously established clinical cardiovascular benefits of eplerenone administration (Pitt et al., 2003), in conjunction with our *in vitro* findings may pave the way for pre-clinical and clinical trials for the treatment of vascular calcification with eplerenone therapy. Inhibiting 11 β -HSD isoenzymes and subsequently diminishing vascular calcification may also represent a novel potential pharmaceutical target for clinical intervention.

Acknowledgements

This work was supported by an Institute Strategic Programme Grant (BB/J004316/1) and Institute Career Path Fellowship (BB/F023928/1) funding from the Biotechnology and Biological Sciences Research Council (BBSRC). The authors are grateful for support from the BHF Centre for Research Excellence.

References

1. Bellows CG, Heersche JN, Aubin JE. Determination of the capacity for proliferation and differentiation of osteoprogenitor cells in the presence and absence of dexamethasone. *Dev Biol.* 1990 140(1):132-8.
2. Block GA. Prevalence and clinical consequences of elevated Ca x P product in hemodialysis patients. *Clin Nephrol.* 2000 54(4):318-24.
3. Boraldi F, Bartolomeo A, Li Q, Uitto J, Quaglini D. Changes in dermal fibroblasts from *Abcc6*^{-/-} mice are present before and after the onset of ectopic tissue mineralization. *J Invest Dermatol.* 2014 134(7):1855-61.
4. Chapman KE, Coutinho AE, Zhang Z, Kipari T, Savill JS, Seckl JR. Changing glucocorticoid action: 11 β -hydroxysteroid dehydrogenase type 1 in acute and chronic inflammation. *J Steroid Biochem Mol Biol.* 2013 Sep;137:82-92. doi: 10.1016/j.jsbmb.2013.02.002.
5. Cheng SL, Yang JW, Rifas L, Zhang SF, Avioli LV. Differentiation of human bone marrow osteogenic stromal cells in vitro: induction of the osteoblast phenotype by dexamethasone. *Endocrinology.* 1994 134(1):277-86.
6. Chmielewski V, Drupt F, Morfin R. Dexamethasone-induced apoptosis of mouse thymocytes: prevention by native 7 α -hydroxysteroids. *Immunol Cell Biol.* 2000 78(3):238-46.
7. Chrysis D, Zaman F, Chagin AS, Takigawa M, Sälvendahl L. Dexamethasone induces apoptosis in proliferative chondrocytes through activation of caspases and suppression of the Akt-phosphatidylinositol 3'-kinase signaling pathway. *Endocrinology.* 2005 146(3):1391-7.

8. Diaz de Barboza G, Guizzardi S, Tolosa de Talamoni N. Molecular aspects of intestinal calcium absorption. *World J Gastroenterol*. 2015 21(23):7142-54. doi: 10.3748/wjg.v21.i23.7142.
9. Guicciardi ME, Mott JL, Bronk SF, Kurita S, Fingas CD, Gores GJ. Cellular inhibitor of apoptosis 1 (cIAP-1) degradation by caspase 8 during TNF-related apoptosis-inducing ligand (TRAIL)-induced apoptosis. *Exp Cell Res*. 2011 317(1):107-16. doi: 10.1016/j.yexcr.2010.10.005.
10. Hadoke PW, Macdonald L, Logie JJ, Small GR, Dover AR, Walker BR. Intravascular glucocorticoid metabolism as a modulator of vascular structure and function. *Cell Mol Life Sci*. 2006 63(5):565-78.
11. Hadoke PW, Kipari T, Seckl JR, Chapman KE. Modulation of 11 β -hydroxysteroid dehydrogenase as a strategy to reduce vascular inflammation. *Curr Atheroscler Rep*. 2013 15(5):320. doi: 10.1007/s11883-013-0320-1.
12. Harris HJ, Kotelevtsev Y, Mullins JJ, Seckl JR, Holmes MC. Intracellular regeneration of glucocorticoids by 11 β -hydroxysteroid dehydrogenase (11 β -HSD)-1 plays a key role in regulation of the hypothalamic-pituitary-adrenal axis: analysis of 11 β -HSD-1-deficient mice. *Endocrinology*. 2001 142(1):114-20.
13. Hartmann K, Koenen M, Schauer S, Wittig-Blaich S, Ahmad M, Baschant U, Tuckermann JP. Molecular Actions of Glucocorticoids in Cartilage and Bone During Health, Disease, and Steroid Therapy. *Physiol Rev*. 2016 96(2):409-47. doi: 10.1152/physrev.00011.2015.
14. Hirata K, Helal F, Hadgraft J, Lane ME. Formulation of carbenoxolone for delivery to the skin. *Int J Pharm*. 2013 20;448(2):360-5. doi: 10.1016/j.ijpharm.2013.03.045.
15. Jaffe IZ, Tintut Y, Newell BG, Demer LL, Mendelsohn ME. Mineralocorticoid receptor activation promotes vascular cell calcification. *Arterioscler Thromb Vasc Biol*. 2007 27(4):799-805.
16. Johnson K, Polewski M, van Etten D, Terkeltaub R. Chondrogenesis mediated by PPI depletion promotes spontaneous aortic calcification in NPP1 $^{-/-}$ mice. *Arterioscler Thromb Vasc Biol*. 2005 25(4):686-91.
17. Johnson KA, Polewski M, Terkeltaub RA. Transglutaminase 2 is central to induction of the arterial calcification program by smooth muscle cells. *Circ Res*. 2008 Mar 14;102(5):529-37. doi: 10.1161/CIRCRESAHA.107.154260.
18. Kirton JP, Wilkinson FL, Canfield AE, Alexander MY. Dexamethasone downregulates calcification-inhibitor molecules and accelerates osteogenic

- differentiation of vascular pericytes: implications for vascular calcification. *Circ Res.* 2006 98(10):1264-72.
19. Kaur K, Hardy R, Ahasan MM, Eijken M, van Leeuwen JP, Filer A, Thomas AM, Raza K, Buckley CD, Stewart PM, Rabbitt EH, Hewison M, Cooper MS. Synergistic induction of local glucocorticoid generation by inflammatory cytokines and glucocorticoids: implications for inflammation associated bone loss. *Ann Rheum Dis.* 2010 69(6):1185-90.
20. Li ZY, Jiang YM, Liu YM, Guo Z, Shen SN, Liu XM, Pan RL. Saikosaponin D acts against corticosterone-induced apoptosis via regulation of mitochondrial GR translocation and a GR-dependent pathway. *Prog Neuropsychopharmacol Biol Psychiatry.* 2014 53:80-9. doi: 10.1016/j.pnpbp.2014.02.010.
21. Liu B, Zhang H, Xu C, Yang G, Tao J, Huang J, Wu J, Duan X, Cao Y, Dong J. Neuroprotective effects of icariin on corticosterone-induced apoptosis in primary cultured rat hippocampal neurons. *Brain Res.* 2011 1375:59-67. doi: 10.1016/j.brainres.2010.12.053
22. Longenecker JP, Kilty LA, Johnson LK. Glucocorticoid inhibition of vascular smooth muscle cell proliferation: influence of homologous extracellular matrix and serum mitogens. *J Cell Biol.* 1984 98(2):534-40.
23. Mackenzie NC, MacRae VE. The role of cellular senescence during vascular calcification: a key paradigm in aging research. *Curr Aging Sci.* 2011 4(2):128-36.
24. Mackenzie NC, Zhu D, Longley L, Patterson CS, Kommareddy S, MacRae VE. MOVAS-1 cell line: a new in vitro model of vascular calcification. *Int J Mol Med.* 2011 27(5):663-8. doi: 10.3892/ijmm.2011.631.
25. Mackenzie NC, Zhu D, Milne EM, van 't Hof R, Martin A, Darryl Quarles L, Millán JL, Farquharson C, MacRae VE. Altered bone development and an increase in FGF-23 expression in *Enpp1(-/-)* mice. *PLoS One.* 2012;7(2):e32177. doi: 10.1371/journal.pone.0032177.
26. Mackenzie NC, Staines KA, Zhu D, Genever P, Macrae VE. miRNA-221 and miRNA-222 synergistically function to promote vascular calcification. *Cell Biochem Funct.* 2014 Mar;32(2):209-16. doi: 10.1002/cbf.3005
27. Molnar GA, Lindschau C, Dubrovskaja G, Mertens PR, Kirsch T, Quinkler M, Gollasch M, Wresche S, Luft FC, Muller DN, Fiebeler A. Glucocorticoid-related signaling effects in vascular smooth muscle cells. *Hypertension.* 2008 May;51(5):1372-8. doi: 10.1161/HYPERTENSIONAHA.107.105718.

28. Montes de Oca A, Guerrero F, Martinez-Moreno JM, Madueño JA, Herencia C, Peralta A, Almaden Y, Lopez I, Aguilera-Tejero E, Gundlach K, Büchel J, Peter ME, Passlick-Deetjen J, Rodriguez M, Muñoz-Castañeda JR. Magnesium inhibits Wnt/ β -catenin activity and reverses the osteogenic transformation of vascular smooth muscle cells. *PLoS One*. 2014 9(2):e89525. doi: 10.1371/journal.pone.0089525.
29. Mori K, Shioi A, Jono S, Nishizawa Y, Morii H. Dexamethasone enhances In vitro vascular calcification by promoting osteoblastic differentiation of vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 1999 19(9):2112-8.
30. Murshed M, Harmey D, Millán JL, McKee MD, Karsenty G. Unique coexpression in osteoblasts of broadly expressed genes accounts for the spatial restriction of ECM mineralization to bone. *Genes Dev*. 2005 19(9):1093-104.
31. Nakagawa Y, Ikeda K, Akakabe Y, Koide M, Uraoka M, Yutaka KT, Kurimoto-Nakano R, Takahashi T, Matoba S, Yamada H, Okigaki M, Matsubara H. Paracrine osteogenic signals via bone morphogenetic protein-2 accelerate the atherosclerotic intimal calcification in vivo. *Arterioscler Thromb Vasc Biol*. 2010 30(10):1908-15. doi: 10.1161/ATVBAHA.110.206185.
32. Pitt, B., Zannad, F., Remme, W. J., Cody, R., Castaigne A, Perez A, Palensky J, Wittes J. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N. Engl. J. Med*. 1999 341(10):709-17.
33. Pitt B, Reichek N, Willenbrock R, Zannad F, Phillips RA, Roniker B, Kleiman J, Krause S, Burns D, Williams GH. Effects of eplerenone, enalapril, and eplerenone/enalapril in patients with essential hypertension and left ventricular hypertrophy: the 4E-left ventricular hypertrophy study. *Circulation*. 2003 108(15):1831-8.
34. Preusch MR, Rattazzi M, Albrecht C, Merle U, Tuckermann J, Schütz G, Blessing E, Zoppellaro G, Pauletto P, Krempien R, Rosenfeld ME, Katus HA, Bea F. Critical role of macrophages in glucocorticoid driven vascular calcification in a mouse-model of atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2008 28(12):2158-64. doi: 10.1161/ATVBAHA.108.174128.
35. Proudfoot D, Skepper JN, Hegyi L, Bennett MR, Shanahan CM, Weissberg PL. Apoptosis regulates human vascular calcification by apoptotic bodies. *Circ Res*. 2000 87:1055–62.
36. Reynolds JL, Skepper JN, McNair R, Kasama T, Gupta K, Weissberg PL, Jahnchen-Dechent W, Shanahan CM. Multifunctional roles for serum protein fetuin-a in

- inhibition of human vascular smooth muscle cell calcification. *J Am Soc Nephrol.* 2005 16(10):2920-30.
37. Richardson RV, Batchen EJ, Denvir MA, Gray GA, Chapman KE. Cardiac GR and MR: From Development to Pathology. *Trends Endocrinol Metab.* 2015 pii: S1043-2760(15)00201-5. doi: 10.1016/j.tem.2015.10.001.
38. Shen CN, Slack JM, Tosh D. Molecular basis of transdifferentiation of pancreas to liver. *Nat Cell Biol.* 2000 2(12):879-87.
39. Shui C, Scutt AM. Mouse embryo-derived NIH3T3 fibroblasts adopt an osteoblast-like phenotype when treated with 1 α ,25-dihydroxyvitamin D(3) and dexamethasone in vitro. *J Cell Physiol.* 2002 193(2):164-72.
40. Skrtic D, Eanes ED. Membrane mediated precipitation of calcium phosphate in model liposomes with matrix vesicle-like lipid composition. *Bone Miner.* 1992 16:109 –119.
41. Son BK, Akishita M, Iijima K, Eto M, Ouchi Y. Mechanism of pi-induced vascular calcification. *J Atheroscler Thromb.* 2008 15(2):63-8.
42. Staines KA, Zhu D, Farquharson C, MacRae VE. Identification of novel regulators of osteoblast matrix mineralization by time series transcriptional profiling. *J Bone Miner Metab.* 2014 32(3):240-51. doi: 10.1007/s00774-013-0493-2.
43. Sun K, Myatt L. Enhancement of glucocorticoid-induced 11 β -hydroxysteroid dehydrogenase type 1 expression by proinflammatory cytokines in cultured human amnion fibroblasts. 2003 144(12):5568-77.
44. Versaci F, Gaspardone A, Tomai F, Ribichini F, Russo P, Proietti I, Ghini AS, Ferrero V, Chiariello L, Giofrè PA, Romeo F, Crea F; Immunosuppressive Therapy for the Prevention of Restenosis after Coronary Artery Stent Implantation Study. *J Am Coll Cardiol.* 2002 40(11):1935-42.
45. Vizzardi E, Nodari S, Caretta G, D'Aloia A, Pezzali N, Faden G, Lombardi C, Raddino R, Metra M, Dei Cas L. Effects of spironolactone on long-term mortality and morbidity in patients with heart failure and mild or no symptoms. *Am J Med Sci.* 2014 347(4):271-6. doi: 10.1097/MAJ.0b013e31829dd6b1.
46. Voelkl J, Alesutan I, Leibrock CB, Quintanilla-Martinez L, Kuhn V, Feger M, Mia S, Ahmed MS, Rosenblatt KP, Kuro-O M, Lang F. Spironolactone ameliorates PIT1-dependent vascular osteoinduction in klotho-hypomorphic mice. *J Clin Invest.* 2013 123(2):812-22. doi: 10.1172/JCI64093
47. Walsh S, Jordan GR, Jefferiss C, Stewart K, Beresford JN. High concentrations of dexamethasone suppress the proliferation but not the differentiation or further

- maturation of human osteoblast precursors in vitro: relevance to glucocorticoid-induced osteoporosis. *Rheumatology (Oxford)*. 2001 40(1):74-83.
48. Wada H. Glucocorticoids: mediators of vertebrate ontogenetic transitions. *Gen Comp Endocrinol*. 2008 156(3):441-53. doi: 10.1016/j.ygcen.2008.02.004.
 49. Zhu D, Mackenzie NC, Millán JL, Farquharson C, MacRae VE. The appearance and modulation of osteocyte marker expression during calcification of vascular smooth muscle cells. *PLoS One*. 2011;6(5):e19595. doi: 10.1371/journal.pone.0019595.
 50. Zhu D, Mackenzie NC, Farquharson C, Macrae VE. Mechanisms and clinical consequences of vascular calcification. *Front Endocrinol (Lausanne)*. 2012 6;3:95. doi: 10.3389/fendo
 51. Zhu D, Mackenzie NC, Millan JL, Farquharson C, MacRae VE. A protective role for FGF-23 in local defence against disrupted arterial wall integrity? *Mol Cell Endocrinol*. 2013 372(1-2):1-11. doi: 10.1016/j.mce.2013.03.008.
 52. Zhu D, Mackenzie NC, Millan JL, Farquharson C, Macrae VE. Upregulation of IGF2 expression during vascular calcification. *J Mol Endocrinol*. 2014 52(2):77-85. doi: 10.1530/JME-13-0136.
 53. Zhu D, Mackenzie NC, Shanahan CM, Shroff RC, Farquharson C, MacRae VE. BMP-9 regulates the osteoblastic differentiation and calcification of vascular smooth muscle cells through an ALK1 mediated pathway. *J Cell Mol Med*. 2015 19(1):165-74. doi: 10.1111/jcmm.12373.

Figure legends

Figure 1: Dexamethasone, corticosterone and 11-DHC all glucocorticoids induce VSMC calcification. Effect of (A) Dexamethasone (1-100nM) (B,C) Corticosterone (white bar) and 11-dehydrocorticosterone (11DHC; filled bar) (1-100nM) in the presence of standard and charcoal-stripped Fetal Bovine Serum (FBS) respectively and (D) Carbenoxolone (10 μ M) in the presence or absence of 11DHC (100 nM) on calcium deposition in VSMCs cultured in high phosphate (Pi) (3mM Pi) for 7 days, as determined by alizarin red staining and/or quantitative HCL leaching (μ g/mg protein) (n=6). Results are presented as mean \pm S.E.M. *P<0.05; **P<0.01; ***P<0.001 compared with corresponding 0 nM treatment.

Figure 2: Endogenous glucocorticoids induce VSMC calcification through the Mineralocorticoid Receptor (MR). Effect of (A) Mifepristone (10 μ M) and (B) Eplerenone (10 μ M) in the presence of Corticosterone (100nM) and (C) Mifepristone (10 μ M) and (D) Eplerenone (10 μ M) in the presence of 11-dehydrocorticosterone (11HDC; 100nM) on calcium deposition in VSMCs cultured in high phosphate (Pi) (3mM Pi) for 7 days, as determined by quantitative HCL leaching (μ g/mg protein) (n=6). Results are presented as mean \pm S.E.M. *P<0.05; **P<0.01; ***P<0.001.

Figure 3: Endogenous glucocorticoids do not induce osteogenic marker expression in VSMCs. Fold change in the mRNA expression of osteogenic markers (A) *Pit-1* (B) *Osx* and (C) *Bmp2* (n=4) following treatment with Corticosterone (white bar) and 11-dehydrocorticosterone (11DHC; filled bar) (1-100nM). 0 nM treatment expressed as 1 to indicate fold change for each gene of interest. Results are presented as mean \pm S.E.M. *P<0.05; **P<0.01; ***P<0.001 compared with corresponding 0 nM treatment.

Figure 4: Endogenous glucocorticoids induce apoptosis in VSMCs. Effect of Corticosterone (1-100nM) on (A) Alamar blue uptake (OD 570nm-620nm) (B) Trypan Blue uptake (%) (C) and (D) Apoptotic nuclei as determined by DAPI staining (%) and (E) Annexin V staining of isolated cells as assessed by FACs analysis. Results are presented as mean \pm S.E.M. *P<0.05; **P<0.01; ***P<0.001 compared with corresponding 0 nM treatment.




Fig. 1




Fig. 2




Fig. 3




Fig. 4